



(02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

POULTRY SUPREME

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2500	2841
Protein (%)	16.50	18.75
Calcium (%)	3.45	3.92
Phosphorus (%)	0.48	0.55
Fibre (%)	5.38	6.11
Fat (%)	5.52	6.27

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Poultry Supreme has been formulated to suit all breeds of layers from point of lay (18 - 20 weeks of age).

Breeding & Show Birds

As a result of a high analysis, Poultry Supreme is not only well suited for those wishing quality egg production & stock health it is also well suited to those wishing to breed and show poultry stock.

- **Proteins** - No urea, all protein in the form of true proteins sources
- **High Energy / Analysis** - For superior egg production and bird health
- **Trace Mineral Premix** - To assist in maintaining nutritional balance and layer health
- **Roxaphyll** - For golden yokes
- **Course Crumble and Short Cut Pellet** - easier acceptance and intake by layers, especially bantams
- **Clearly Identifiable Bags** - Avoids using the wrong product

Other Specialty Products

- Chicken Starter
- Free Range Layer
- Broiler Diets
- Pullet Grower
- Free Range Scratch Mix
- Game Bird Diets
- Back Yard Layer
- Turkey Diets
- Duck Diets

General Information

- Introduce to birds from 18 - 20 weeks of age
- Intake varies from 105 grams - 140 grams per day depending on bird health & breed, lighting, housing, climate conditions and water supply
- Feeding hens to production is important to maximise bird health, egg production and yoke colour
- Fresh, clean water to be accessible at all times
- Greens can be fed in conjunction with pellets
- After 52 weeks of age a non commercial layer pellet can be used to match layers egg production
- Older hens (>2 years) egg production will decline rapidly

- *Egg production rates will vary according to a range of external factors including bird age, housing, lighting, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draft free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers need to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick layers to prevent possible outbreaks*
- *Dead layers need to be removed straight away and disposed of.*

Revision: 20071013