



(02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

DUCK GROWER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg ¹	2500	2808
Protein (%)	17.00	19.10
Calcium (%)	2.00	2.24
Phosphorus (%)	0.50	0.56
Fibre (%)	5.70	6.40
Fat (%)	4.55	5.11

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Duck Grower has been formulated to suit all breeds of ducks from 4 - 5 weeks of age to desired weight or for breeding for domestic use.

Chicken Starter suitable as a starter diet (0 - 4 weeks or age).

- **High Quality Proteins** - No urea, all protein in the form of true proteins sources
- **Specialty Product** - Formulated especially for domestic ducks for both meat and breeding
- **Energy** - To assist with growth
- **Trace Mineral Premix** - To assist in maintaining nutritional balance and bird health
- **Pellets** - easier acceptance and intake by ducks
- **Clearly Identifiable Bags** - Avoids using the wrong product

Other Specialty Products

- Chick Starter
- Free Range Layer
- Turkey Diets
- Pullet Grower
- Commercial Layer
- Game Bird Diets
- Back Yard Layer
- Free Range Scratch Mix
- Broiler Diets

General Information

- A complete pellet (diet) providing all requirements for growing ducks
- Suitable to be fed to all breeds of ducks from 4 - 5 weeks of age to desired weight or breeding
- The diet should be fed ad-lib to all ducks, no other supplement is required
- Fresh, clean water to be accessible at all times close to feeders
- Feed intake will vary at times, due to climate and general housing of young ducks
- Average feed consumption from 4 weeks to 8 weeks 6.00 - 6.50 kg

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, disease, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draught free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick birds to prevent possible outbreaks*
- *Dead birds need to be removed straight away and disposed of.*

Revision: 20070808