



(02) 9625 2311
 (02) 9832 2469
 www.thevellagroup.com.au
 sales@thevellagroup.com.au
 PO Box 73, Doonside. NSW. 2767
 96 Glendenning Rd Plumpton. NSW. 2761

PULLET GROWER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2300	2614
Protein (%)	16.00	18.18
Calcium (%)	1.31	1.9
Phosphorus (%)	0.48	0.55
Fibre (%)	6.70	7.61
Fat (%)	3.84	4.36

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

Pullet Grower has been formulated to suit all breeds of pullets from 6 - 7 weeks to point of lay (18 - 20 weeks of age).

This is a complete feed and no other supplement is required

- **High Quality Proteins** - No urea, all protein in the form of true proteins sources
- **Coccidiostat** - Assist chicks against coccidiosis
- **Energy** - To assist in healthy growth
- **Trace Mineral Premix** - To assist in maintaining nutritional balance and pullet health
- **Course Crumble** - assist in transition to layer pellets also suited to Bantams.
- **Clearly Identifiable Bags** - Avoids using the wrong product
- **Easy Lift Bag** - For your safety

Other Specialty Products

- Chicken Starter
- Commercial Layer
- Broiler Diets
- Back Yard Layer
- Free Range Scratch Mix
- Game Bird Diets
- Free Range Layer
- Turkey Diets
- Duck Diets

General Information

- A complete diet available in a course crumble form, providing important nutrients for growing pullets
- Suitable to be fed to all breeds of birds from 6 weeks to point of lay (18—20 weeks)
- A coccidiostat has been added to assist in the control of coccidiosis
- Fresh, clean water to be accessible at all times
- Feed intake will vary at times, due to climate and general housing of pullets
- As pullets develop (age) feed intake will increase rapidly until maturity
- Average consumption from 6 weeks of age to point of lay, approximately 5—7 kg of feed per bird

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draft free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick pullets to prevent possible outbreaks*
- *Dead pullets need to be removed straight away and disposed of.*

Revision: 20070808