

## CHICKEN STARTER

Typical Analysis	As Fed	Dry Matter
Energy KCAL / kg	2700	3000
Protein (min) (%)	20.00	22.20
Calcium (min) (%)	1.00	1.14
Phos (min) (%)	0.61	0.69
Fibre (max) (%)	6.00	6.60
Fat (max) (%)	6.00	6.60

TRACE MINERAL PREMIX ADDED, FOR MORE DETAILED ANALYSIS PLEASE CONTACT VELLA STOCK FEEDS.

**Chicken Starter** has been formulated to suit all breeds of layer chickens from day old to 6 - 7 weeks of age.

This is a complete feed and no other supplement is required.

Contains a natural herbal coccidiostat.

- **Probiotic and Prebiotic** - Good bacteria for gut health
- **Coccidiostat** - Regano, a natural herbal coccidiostat used, no withholding period for meat and or egg consumption. Also suitable for ducklings to consume.
- **Natural Betaine** - Support chicks during times of heat stress
- **Vegetable Proteins** - No restricted animal materials (RAM) used in our diets
- **Vitamins** - Species specific vitamins & minerals included to support health and production
- **Antioxidant** - Assist health
- **Ducklings** - Suitable for Ducklings
- **No** added hormones or antibiotics used

### Other Specialty Products

- Pullet Grower
- Poultry Supreme
- Meat Bird Finisher / Breeder
- Back Yard Layer
- Free Range Scratch Mix
- Commercial Diets
- Free Range Layer
- Meat Bird Starter
- Duck Grower

### General Information

- A complete crumble (diet) providing all requirements for young chickens
- Suitable to be fed to all breeds of chickens from day old to 6 - 7 weeks of age
- The diet should be fed ad-lib to all chickens, no other supplement is required
- A coccidiostat has been added to assist in the control of coccidiosis
- Fresh, clean water to be accessible at all times (24 /7)
- To assist young birds in the first few days of life, change feed regularly
- Feed intake will vary at times, due to climate and general housing of chickens
- Average consumption from 0 - 6 / 7 weeks of age, approximately 2.0 - 2.5 kg of feed per chicken

- *Growth rates will vary according to a range of external factors including housing, variable weather conditions, predators, inoculations and hygiene to name a few*
- *Phase supplementary feeding both in and out gradually to help avoid set backs.*
- *Housing should be well vented but draft free*
- *Ensure clean, cool water is accessible at all times*
- *Feeders and waterers needs to be easily accessible and cleaned of litter on a regular basis*
- *Quarantine sick chicks to prevent possible outbreaks*
- *Dead chicks need to be removed straight away and disposed of.*

Revision:20170906